



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Skin test reactivity to hymenoptera venom after venom immunotherapy correlates inversely with the IgG/IgE ratio

Saulite, Ieva ; Hoetzenecker, Wolfram ; Guenova, Emmanuella ; Schmid-Grendelmeier, Peter ; Glatz, Martin

Abstract: BACKGROUND: Skin test reactivity to hymenoptera venom and venom-specific IgE are important for diagnosing venom allergy and deciding on the appropriate allergen for venom immunotherapy (VIT). Longitudinal data on skin test reactivity during VIT and their correlation with venom-specific immunoglobulin (Ig)E and IgG are scarce. METHODS: We retrospectively analyzed shifts in skin test reactivity and serum levels of venom-specific IgE and IgG in patients allergic to hymenoptera venom before the initiation of VIT with ultrarush therapy and after 3 years of VIT. RESULTS: Fifty-four patients received ultrarush desensitization and subsequent VIT with wasp venom, 26 with honeybee venom, and 8 with both wasp and honeybee venom. Hymenoptera-specific skin test reactivity decreased during VIT in most patients, and became negative in 8% of the wasp-allergic patients and in 25% of the honeybee-allergic patients. Serum levels of venom-specific IgE positively correlated to skin test reactivity before VIT, but did not change significantly during VIT. IgG serum levels and the IgG/IgE ratio increased during VIT in most patients. A high IgG/IgE ratio correlated with low skin test reactivity after 3 years of VIT. CONCLUSIONS: The correlation between a high venom-specific IgG/IgE ratio and low skin test reactivity after VIT may be interesting for future investigations that assess its role as a potential marker for VIT efficacy.

DOI: <https://doi.org/10.1159/000481255>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-143164>

Journal Article

Published Version

Originally published at:

Saulite, Ieva; Hoetzenecker, Wolfram; Guenova, Emmanuella; Schmid-Grendelmeier, Peter; Glatz, Martin (2017). Skin test reactivity to hymenoptera venom after venom immunotherapy correlates inversely with the IgG/IgE ratio. *International Archives of Allergy and Immunology*, 174(3-4):190-199.

DOI: <https://doi.org/10.1159/000481255>

Skin Test Reactivity to Hymenoptera Venom after Venom Immunotherapy Correlates Inversely with the IgG/IgE Ratio

Ieva Saulite^{a–c} Wolfram Hoetzenecker^{a, b} Emmanuella Guenova^{a, b}
Peter Schmid-Grendelmeier^a Martin Glatz^a

^aAllergy Unit, Department of Dermatology, University and University Hospital of Zurich, Zurich, and ^bDepartment of Dermatology/Allergology, Cantonal Hospital St. Gallen, St. Gallen, Switzerland; ^cDepartment of Infectiology and Dermatology, Riga Stradins University, Riga, Latvia

Keywords

Hymenoptera venom allergy · Immunoglobulin E · Immunoglobulin G · Skin test · Venom immunotherapy

Abstract

Background: Skin test reactivity to hymenoptera venom and venom-specific IgE are important for diagnosing venom allergy and deciding on the appropriate allergen for venom immunotherapy (VIT). Longitudinal data on skin test reactivity during VIT and their correlation with venom-specific immunoglobulin (Ig)E and IgG are scarce. **Methods:** We retrospectively analyzed shifts in skin test reactivity and serum levels of venom-specific IgE and IgG in patients allergic to hymenoptera venom before the initiation of VIT with ultrarush therapy and after ≥ 3 years of VIT. **Results:** Fifty-four patients received ultrarush desensitization and subsequent VIT with wasp venom, 26 with honeybee venom, and 8 with both wasp and honeybee venom. Hymenoptera-specific skin test reactivity decreased during VIT in most patients, and became negative in 8% of the wasp-allergic patients and in 25% of the honeybee-allergic patients. Serum levels of venom-specific IgE positively correlated to skin test reactivity before VIT, but did not change significantly during VIT.

IgG serum levels and the IgG/IgE ratio increased during VIT in most patients. A high IgG/IgE ratio correlated with low skin test reactivity after ≥ 3 years of VIT. **Conclusions:** The correlation between a high venom-specific IgG/IgE ratio and low skin test reactivity after VIT may be interesting for future investigations that assess its role as a potential marker for VIT efficacy.

© 2017 S. Karger AG, Basel

Introduction

Wasp or honeybee venom allergy affects 0.3–7.5% of the general population [1]. Venom immunotherapy (VIT) is usually performed in individuals experiencing severe anaphylactic reactions [2]. VIT comprises venom injections during an up dosing phase over several weeks and a maintenance phase for 3–5 years. Recently, ultrarush therapy protocols for up dosing during only a few hours have been introduced to clinical routine. Ultrarush

This study was carried out at the Allergy Unit, Department of Dermatology, University and University Hospital of Zurich, Zurich, Switzerland

therapy is safe, fast, provides optimal compliance, and is thus cost effective [3]. The accurate diagnosis of venom allergy and the identification of the causative insect are decisive in selecting the appropriate venom to use in VIT. The diagnostic workup comprises a thorough history, skin tests, and in vitro tests [4]. For skin testing, increasing concentrations of venom are injected intracutaneously until a positive reaction with a wheal and surrounding erythema (flare) is observed [5]. The skin test is considered the gold standard for the diagnosis of hymenoptera venom allergy, because it is sensitive to distinguishing sensitization to different types of venom, gives results within a few minutes, and may simulate the real-life situation of a sting better than in vitro tests [2, 4]. In vitro tests usually comprise the detection of immunoglobulin E (IgE) to whole venom and/or venom components such as the major wasp allergens Ves v 1 and Ves v 5 or the major bee allergen Api m 1, well as the detection of venom-specific IgG [6–8].

However, despite the regular use of skin tests and the measurement of venom-specific IgE and IgG before VIT, little is known about the longitudinal shifts of these parameters and the correlations between them during VIT. This is particularly true for patients who started VIT with ultrarush up dosing [3]. We therefore carried out this study on wasp-allergic and/or honeybee venom-allergic patients undergoing VIT after ultrarush up dosing, to assess (i) longitudinal shifts of skin test reactivity and venom-specific antibodies, and (ii) correlations between skin test reactivity and venom-specific antibodies before and after ≥ 3 years of VIT.

Methods

Patients

This was a retrospective, single-center study. We screened all medical files of patients undergoing ultrarush therapy and consecutive VIT with wasp or/and honeybee venom between 2010 and 2012 at the Allergy Unit, Department of Dermatology, University Hospital of Zurich. Inclusion criteria were (i) diagnosis of wasp and/or bee venom allergy that qualified for VIT, (ii) an age of ≥ 18 years, (iii) ultrarush therapy and VIT for a minimum of 3 years according to the standard protocol of our department (see below), and (iv) data on venom skin test reactivity and venom-specific IgE and IgG from before ultrarush therapy versus after a minimum of 3 years of VIT. Exclusion criteria were (i) previous VIT with wasp and/or bee venom, (ii) the patient being a beekeeper, and (iii) systemic mastocytosis. The protocol for ultrarush sensitization and the maintenance phase of VIT was as follows. During ultrarush therapy, patients received increasing concentrations of total 111 μg of aqueous wasp or bee venom (Pharmalgen, ALK-Abello, Hørsholm, Denmark) into the dorsal upper arm within 2.5–3.5 h [3].

During the maintenance phase of VIT, patients received 100 μg of aluminium-adsorbed depot venom extract (Alutard SQ, ALK-Abello) every 4 weeks during the first year, every 5 weeks during the second year, and every 6 weeks during the third year after ultrarush therapy. Injections during the maintenance phase were either applied in our department or with a health care provider. Injection protocols of the maintenance phase were checked for violations of our standard protocol during the follow-up visit.

The study was conducted in accordance with the Declaration of Helsinki. For all patients, informed consent forms for the study were obtained in accordance with the Biobank Project (EK No. 647).

Clinical and Laboratory Parameters

In patients fulfilling the inclusion criteria, the following data were retrieved from the medical files: age; gender; a history of atopy defined as allergic rhinoconjunctivitis, and/or allergic asthma, or atopic dermatitis; the severity of the allergic reaction to the wasp or bee sting as graded by the criteria of Mueller [9]; the duration of VIT; the serum tryptase level; serum levels of IgE specific to whole wasp or bee venom (Ves v 1, Ves v 5, and Api m 1) before and after VIT; serum levels of wasp or bee venom-specific IgG before and after VIT; skin test reactivity to wasp or bee venom before and after VIT. For the diagnostic workup, laboratory tests and skin tests before VIT were performed >8 weeks after the culprit wasp or bee sting. At follow-up during or after VIT, laboratory tests and skin tests were performed ≥ 3 weeks after the last venom injection.

The levels of venom-specific IgE were determined by ImmunoCAP (Phadia, Thermo Fisher Scientific, Uppsala, Sweden), and those of venom-specific IgG were determined by a cellular allergy activation test followed by an enzyme-linked immunosorbent assay (Bühlmann Laboratories AG, Schönenbuch, Switzerland). For skin tests, 0.1 mL of increasing concentrations of aqueous extracts of wasp or bee venom (Pharmalgen) was injected intradermally into the lower forearms of patients. The 3 lowest venom concentrations (0.00001, 0.001, and 0.01 $\mu\text{g}/\text{mL}$), the negative and positive controls, were administered simultaneously. If these venom concentrations were negative, the 2 highest venom concentrations (0.1 and 1.0 $\mu\text{g}/\text{mL}$) were consecutively injected intradermally after 15 min each. All patients were allocated to 1 of 3 groups: group 1, patients receiving ultrarush therapy and VIT exclusively with wasp venom; group 2, patients receiving ultrarush therapy and VIT exclusively with honeybee venom; and group 3, patients receiving ultrarush therapy and VIT with both wasp and honeybee venom.

Statistics

Statistical analysis was performed with R software v3.2.3 [10]. The distribution of continuous data was tested for normality by means of the Kolmogorov-Smirnov and Lilliefors tests. Outliers of data points were visualized with Q-Q plots. We assumed a non-normal distribution of continuous data among all groups and used nonparametric tests for further statistical analysis (the Wilcoxon rank test and the Kruskal-Wallis test). Frequencies of positive skin test results before versus after ≥ 3 years of VIT initiated by ultrarush were compared using the χ^2 test with the Yate continuity correction. The dilution series of 0.00001:1 $\mu\text{g}/\text{mL}$ wasp or bee venom to determine skin test reactivity comprised 5 discrete data points, with a 6th data point for negative test results (equal to 10 $\mu\text{g}/\text{mL}$). This series of discrete data points was treated as continuous data for statistical analysis [11]. Continuous data (skin test reactivity




Fig. 1. Patient selection according to inclusion and exclusion criteria. VIT, venom immunotherapy.

and serum antibody titers) before versus after ≥ 3 years of VIT initiated by ultrarush desensitization were therefore compared using the Wilcoxon rank test. Clinical and laboratory metadata (gender, age, anaphylaxis grade, and serum tryptase level) were compared using the χ^2 test or the Kruskal-Wallis test. All p values were adjusted for multiple comparisons with the Benjamini-Hochberg correction [12]. The correlation between skin test reactivity and laboratory and clinical parameters was analyzed using the Spearman rho. Statistical significance for all tests was ascribed to 2-sided α level of the adjusted p values <0.05 , and by 95% confidence intervals for continuous data.


Results

Patients

Between 2010 and 2012, a total of 340 patients received ultrarush therapy and VIT with wasp venom and/or honeybee venom in our department. Eighty-eight patients

(25.8%) fulfilled the inclusion criteria and were further analyzed (Fig. 1). Fifty-four patients (61%) received ultrarush therapy and VIT with wasp venom (“wasp venom group,” 27 females (50%); median age 44 years, interquartile range [IQR] 38–56 years), 26 patients (30%) with honeybee venom (“bee venom group,” 13 females (50%); median age 46 years, IQR 24–52 years), and 8 patients (9%) with both wasp and honeybee venom (“wasp/bee venom group,” 1 female (12.5%); median age 30 years, IQR 26–46 years). There was no statistically significant difference regarding the sex ($p = 0.11$) or age of patients ($p = 0.28$) in the 3 groups. All patients had a history of a systemic allergic reaction of grade II–IV after a wasp or bee sting (median grade III, IQR III–IV) according to Mueller [9]. The median time period between the initiation of VIT (including ultrarush therapy) and the follow-up control visit for laboratory analysis and skin test was 3 years (IQR 3–5 years) in all patient groups.

Fig. 2. Concentration of the respective venom that was necessary to elicit a positive intradermal skin test before and after ≥ 3 years of venom immunotherapy (VIT) with wasp venom (**a**) and bee venom (**b**). The diamond indicates the median venom concentration among all patients. *** $p < 0.001$.



Longitudinal Shifts in Skin Test Reactivity

All patients received a skin test with both wasp and honeybee venom before ultrarush therapy. The majority of patients had a positive skin test before VIT. Only 1 patient in the wasp venom group (2%) and 2 patients in the bee venom group (8%) had a negative skin test before VIT. These 3 patients had an unambiguous history of allergic insect sting reactions and elevated venom-specific IgE to support the diagnosis of hymenoptera venom allergy.

After ≥ 3 years of VIT, a skin test was done on 24 patients in the wasp venom group (44%), 12 in the bee venom group (46%), and 5 in the wasp/bee venom group (63.5%). The skin test was performed at a median of 3 weeks after the previous venom injection (IQR 3–4 weeks). Although most patients remained positive at the skin test, we observed an average decrease of reactivity of 1 venom dilution step in the wasp venom and bee venom groups before versus after ≥ 3 years of VIT (Fig. 2). Only a few patients became negative at the skin test after ≥ 3 years of VIT (Table 1). This decrease was statistically significant in the wasp venom group ($p < 0.001$; 95% CI 1–2), but not in the bee venom group ($p = 0.2$; 95% CI –2 to 0.00002), presumably owing to the smaller number of patients in this group that had a skin test after ≥ 3 years of VIT ($n = 12$). We did not observe a significant shift in skin test reactivity before versus ≥ 3 years of VIT in the wasp/

Table 1. Dynamics of skin test reactivity after ≥ 3 years of venom immunotherapy

	Wasp venom group ($n = 24$)	Bee venom group ($n = 12$)	Wasp/bee venom group ($n = 5$)	
			wasp venom	bee venom
Decrease	17 (71)	5 (42)	2 (60)	3 (40)
Negative	2 (8)	3 (25)	0	0
Unchanged	3 (13)	3 (25)	1 (20)	2 (40)
Increase	2 (8)	1 (8)	1 (20)	1 (20)

Values are expressed as n (%). Patients in all 3 groups underwent skin testing before ultrarush therapy and after ≥ 3 years of venom immunotherapy.

bee venom group, but the number of patients who received a skin test after ≥ 3 years of VIT in this group was too small ($n = 5$) for the statistical analysis.

Laboratory Parameters and Correlation with Skin Test Reactivity

Venom-specific IgE did not change significantly before versus after ≥ 3 years of VIT, in the wasp venom group (IgE to whole wasp venom, $p = 0.84$; Ves v 1, $p = 0.95$; Ves v 5, $p = 0.054$) or in the bee venom group (IgE




Fig. 3. Correlation between the venom concentration necessary to elicit a positive intradermal skin test and the serum levels of IgE to the respective whole venom before and after ≥3 years of venom immunotherapy (VIT) with wasp venom (a) and bee venom (b). The correlation was analyzed using the Spearman rho.




Fig. 4. Serum levels of venom-specific IgG before and after ≥ 3 years of venom immunotherapy (VIT). * $p < 0.05$; *** $p < 0.001$.

to whole bee venom, $p = 1$; Api m 1, $p = 0.78$). Patients with higher serum IgE levels to the whole venom extract had a positive skin test to lower concentrations of the respective venom before and after ≥ 3 years of VIT. This correlation was generally weaker in the wasp venom group than in the bee venom group (Fig. 3). The number of patients in the wasp/bee venom group was too small for statistical analysis.

Serum levels of venom-specific IgG increased before versus after ≥ 3 years of VIT in all groups. This increase was statistically significant in the wasp venom-allergic patients of both the wasp venom group ($p < 0.001$) and the wasp/bee venom group ($p = 0.036$), and in the bee venom-allergic patients of the wasp/bee venom group ($p = 0.02$) but not the bee venom group ($p = 0.3$) (Fig. 4). Unlike with venom-specific IgE, serum levels of venom-specific IgG did not correlate with skin test reactivity to the respective venom at all (data not shown).

The ratio of venom-specific IgG/whole venom-specific IgE (IgG/IgE) significantly increased in the wasp venom group before versus after ≥ 3 years of VIT ($p < 0.01$). In the bee venom group, the IgG/IgE ratio increased in some patients after ≥ 3 years of VIT, but remained unchanged in others (Fig. 5). Patients with a higher venom-specific IgG/IgE ratio had lower skin test reactivity after ≥ 3 years of VIT than patients with a lower IgG/IgE ratio (Fig. 5).

We also investigated the correlation of the metadata, i.e., gender, age, anaphylaxis grade, presence of atopy, and serum tryptase level, with shifts in skin test reactivity and also venom-specific IgE and IgG, but no statistically significant correlation was observed.

Patients with Sting Challenges after VIT

Two bee-allergic patients underwent bee-sting provocation after the completion of VIT. This number was too small for statistical analysis. However, both patients tolerated the sting challenge without systemic or major local reactions, which indicates bee venom tolerance.

Discussion

This was a retrospective study of 88 patients undergoing ultrarush therapy and VIT with wasp and/or honey-bee venom. We analyzed longitudinal shifts in skin test reactivity, serum levels of venom-specific IgE or IgG, and the correlation between these parameters. We found that skin test reactivity usually declined after ≥ 3 years of VIT, and became negative in 8 and 25% of wasp- and bee-allergic patients, respectively. This decline of skin test reactivity during VIT comprised a median 1 dilution of venom in both the wasp venom and bee venom groups. The statistically nonsignificant decline in the bee venom group was presumably due to the smaller number of patients that received a skin test at follow-up ($n = 12$) than there were in the wasp venom group ($n = 24$), and could therefore be a statistical phenomenon.

The decline of skin test reactivity observed in this study is in line with previous findings. In 2 studies on >580 wasp- or bee-allergic patients, skin test reactivity declined in about two-thirds of the wasp-allergic patients and in about half of the bee-allergic patients [13, 14]. There was a wide range of patients in which the skin test




Fig. 5. a Ratio of venom-specific IgG/whole venom-specific IgE before and after ≥ 3 years of venom immunotherapy (VIT). **b** Correlation between the venom concentration necessary to elicit a positive intradermal skin test and the ratio of venom-specific IgG/whole venom-specific IgE after ≥ 3 years of VIT. The correlation was analyzed using the Spearman rho. ** $p < 0.01$.

became negative after VIT, i.e., 6–27% of the wasp-allergic and 2.5–29% of the bee-allergic patients [13–17]. The rate of negative skin tests after VIT appears to be higher in studies that have applied an ultrarush protocol for venom up dosing [16, 17]; this includes our study. It might

therefore be speculated that the higher rate of skin test-negative patients after VIT in our study relates to the use of an ultrarush protocol for venom up dosing. Another explanation could be the use of an aluminium-adsorbed depot extract during the maintenance phase of VIT in our

study. In contrast, previous studies with lower rates of negative skin test reactivity after VIT exclusively used an aqueous extract [13, 14]. However, until now, no differences have been noted in the efficacy of VIT when using these 2 venom extract types, as shown in a study on 27 patients undergoing VIT with a depot extract versus 18 patients undergoing VIT with an aqueous extract [18].

It remains unclear if a decrease in skin test reactivity after VIT indicates protection from allergic reactions at future allergen exposure. The wheal of a positive skin test reaction is caused by local, IgE-mediated mast cell activation, indicates allergen sensitization, and may also hint at a possibly systemic mast cell reaction after allergen exposure, e.g., after a wasp or bee sting [19]. Therefore, some authors attribute a decreased reactivity in these skin tests as a possible indicator of the lower susceptibility of mast cells to an allergen, and hence a hint of an efficacious VIT. They suggest skin test negativity as the goal of a successful VIT [13, 20, 21]. However, only a minority of patients achieve negativity in the skin test after VIT [13, 14, 17], and a decrease in skin test reactivity may not necessarily indicate tolerance of restings after VIT, e.g., at sting challenges [13, 15, 20].

Another possible parameter to assess venom tolerance is a sting challenge, but a tolerated sting challenge does not necessarily predict tolerance of future sting in an individual patient [1, 4, 22–24]. Moreover, up to 17% of patients experience systemic reactions during these sting challenges, even after 1–5 years of VIT [25, 26]. These reactions can be near-fatal or even fatal, experienced in our department (unpubl. data) and also reported by other study groups [20, 27]. This corroborates the need for a reliable and safe parameter to assess VIT efficacy. To support the search for a new parameter that fulfills these requirements, we were interested to further assess possible shifts in laboratory parameters before versus after VIT, and to correlate these shifts with skin test reactivity.

We did not observe a significant shift of venom-specific IgE, but there was an increase in venom-specific IgG before versus after ≥ 3 years of VIT. Some previous studies described the opposite, namely, a significant decrease in venom-specific IgE but unchanged serum levels of venom-specific IgG. This may be attributed to (i) the use of different detection methods such as a radioallergosorbent assay for IgE and another ELISA for IgG [13] compared to the ImmunoCAP and CAST/ELISA used in our study; (ii) a shorter duration of VIT, like 1 year [14] compared to the ≥ 3 years in our study; or (iii) the use of different immunotherapy protocols. Increased IgG antibody levels during VIT have been described pre-

viously, e.g., in a case report on a beekeeper's wife that changed VIT from whole bee extract to pure bee venom [28], and in a study on 40 patients with yellow-jacket venom allergy [29]. However, these studies either described a single patient [28], or the IgG levels were measured already after a few weeks after the onset of VIT and by a method different from ours [29]. A more recent study by Schiavino et al. [17] applied an ultrarush protocol similar to ours, and it described an increase of IgG but unchanged IgE during VIT, as described in our study. It therefore remains possible that the venom up-dosing with ultrarush therapy modulates the immune response to an allergen differently from a conventional up-dosing protocol [30], but the exact mechanisms remain to be elucidated. Ambiguous results regarding shifts of IgG before and after VIT could also be explained by the detection of total IgG versus IgG₄ in different studies. In our study, we measured total IgG rather than IgG₄ only, because IgG₁ also has important protective effects after VIT by blocking IgE-mediated reactions via different mechanisms [31]. For example, IgG can directly bind to an allergen and therefore block the allergen recognition by IgE. IgG also binds the inhibitory FcεRIIb receptor on mast cells and abrogates mast cell activation by IgE [32]. However, a correlation between increased IgG after VIT and protection from allergic reactions after future stings has not yet been proven [33]. This might be due to the possibility that not all IgGs bind to epitopes on allergens that are recognized by IgE, hence allergen recognition by IgE is not totally blocked despite high IgG levels [34]. Therefore, the IgG/IgE ratio might be a better indicator for the protection against allergic reactions [35]. We found an increase of the venom-specific IgG/IgE ratio in all wasp-allergic and in some bee-allergic patients before versus after ≥ 3 years of VIT, which is in line with previous studies [21, 36]. We also investigated the correlation between skin test reactivity and IgG/IgE ratio after 3 years VIT which, to our knowledge, has not yet been published. We found that wasp- and bee-allergic patients with a high IgG/IgE ratio have low skin test reactivity after ≥ 3 years of VIT. But it is unclear if a high IgG/IgE ratio after VIT indeed indicates protection from allergic reactions to wasp or bee restings. Early studies on 176 wasp- or bee-allergic patients did not observe a difference in the outcome of diagnostic sting challenge, depending on the serum levels of venom-specific IgE and IgG or the IgG/IgE ratio [37, 38]. Notably, these were diagnostic studies and they measured the serologic parameters before VIT for diagnostic purposes but not after VIT, as in our study.

Acknowledgements

This study was supported by a Clinical Fellowship of the European Academy of Allergy and Clinical Immunology (2015) (to I.S.), the Swiss National Science Foundation (PMPDP3_151326 to E.G.), the Helmut Horten Foundation (to E.G.), and the Hochspezialisierte Medizin Schwerpunkt Immunologie (HSM-2-Immunologie) Schweiz.

Disclosure Statement

The authors declare no conflict of interest for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Bilo BM, Rueff F, Mosbech H, Bonifazi F, Oude-Elberink JN: Diagnosis of hymenoptera venom allergy. *Allergy* 2005;60:1339–1349.
- Ludman SW, Boyle RJ: Stinging insect allergy: current perspectives on venom immunotherapy. *J Asthma Allergy* 2015;8:75–86.
- Roll A, Hofbauer G, Ballmer-Weber BK, Schmid-Grendelmeier P: Safety of specific immunotherapy using a four-hour ultra-rush induction scheme in bee and wasp allergy. *J Investig Allergol Clin Immunol* 2006;16:79–85.
- Krishna MT, Ewan PW, Diwakar L, Durham SR, Frew AJ, Leech SC, Nasser SM: Diagnosis and management of hymenoptera venom allergy: British Society for Allergy and Clinical Immunology (BSACI) guidelines. *Clin Exp Allergy* 2011;41:1201–1220.
- Strohmeier B, Aberer W, Bokanovic D, Komericki P, Sturm GJ: Simultaneous intradermal testing with hymenoptera venoms is safe and more efficient than sequential testing. *Allergy* 2013;68:542–544.
- Korosec P, Valenta R, Mittermann I, Celesnik N, Erzen R, Zidarn M, Kosnik M: Low sensitivity of commercially available rApi m 1 for diagnosis of honeybee venom allergy. *J Allergy Clin Immunol* 2011;128:671–673.
- Muller UR, Johansen N, Petersen AB, Fromberg-Nielsen J, Haeberli G: Hymenoptera venom allergy: analysis of double positivity to honey bee and *Vespula* venom by estimation of IgE antibodies to species-specific major allergens Api m1 and Ves v5. *Allergy* 2009;64:543–548.
- Manso EC, Morato-Castro FF, Yee CJ, Croce M, Palma MS, Croce J: Honeybee venom specific IgG subclass antibodies in Brazilian beekeepers and in patients allergic to bee stings. *J Investig Allergol Clin Immunol* 1998;8:46–51.
- Mueller HL: Diagnosis and treatment of insect sensitivity. *J Asthma Res* 1966;3:331–333.
- R Core: R: a language and environment for statistical computing. Vienna, 2013. <http://www.r-project.org/>.
- Nevill AM, Atkinson G, Hughes MD, Cooper SM: Statistical methods for analysing discrete and categorical data recorded in performance analysis. *J Sports Sci* 2002;20:829–844.
- Benjamini Y, Hochberg Y: Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc B (Methodological)* 1995;57:289–300.
- Lerch E, Muller UR: Long-term protection after stopping venom immunotherapy: results of re-stings in 200 patients. *J Allergy Clin Immunol* 1998;101:606–612.
- Carballada F, Boquete M, Nunez R, Lombardero M, de la Torre F: Follow-up of venom immunotherapy (VIT) based on conventional techniques and monitoring of immunoglobulin E to individual venom allergens. *J Investig Allergol Clin Immunol* 2010;20:506–513.
- Stoevesandt J, Hofmann B, Hain J, Kerstan A, Trautmann A: Single venom-based immunotherapy effectively protects patients with double positive tests to honey bee and *Vespula* venom. *Allergy Asthma Clin Immunol* 2013;9:33.
- Thurnheer U, Muller U, Stoller R, Lanner A, Hoigne R: Venom immunotherapy in hymenoptera sting allergy. Comparison of rush and conventional hyposensitization and observations during long-term treatment. *Allergy* 1983;38:465–475.
- Schiavino D, Nucera E, Pollastrini E, De Pasquale T, Buonomo A, Bartolozzi F, Lombardo C, Roncallo C, Patriarca G: Specific ultrarush desensitization in hymenoptera venom-allergic patients. *Ann Allergy Asthma Immunol* 2004;92:409–413.
- Cadario G, Marengo F, Raghino E, Rossi R, Gatti B, Cantone R, Bona F, Pellegrino R, Feyles G, Puccinelli P, Burastero SE: Higher frequency of early local side effects with aqueous versus depot immunotherapy for hymenoptera venom allergy. *J Investig Allergol Clin Immunol* 2004;14:127–133.
- Li JT: Allergy testing. *Am Fam Physician* 2002;66:621–624.
- Golden DB: Fatal insect allergy after discontinuation of venom immunotherapy. *J Allergy Clin Immunol* 2001;107:925–926.
- Ewan PW: New insight into immunological mechanisms of venom immunotherapy. *Curr Opin Allergy Clin Immunol* 2001;1:367–374.
- Muller UR, Ring J: When can immunotherapy for insect sting allergy be stopped? *J Allergy Clin Immunol* 2015;3:324–328; quiz 329–330.
- Yunginger JW: The sting – revisited. *J Allergy Clin Immunol* 1979;64:1–2.
- Rueff F, Przybilla B, Muller U, Mosbech H: The sting challenge test in hymenoptera venom allergy. Position Paper of the Subcommittee on Insect Venom Allergy of the European Academy of Allergology and Clinical Immunology. *Allergy* 1996;51:216–225.
- Golden DB, Kwiterovich KA, Kagey-Sobotka A, Valentine MD, Lichtenstein LM: Discontinuing venom immunotherapy: outcome after five years. *J Allergy Clin Immunol* 1996;97:579–587.
- Muller U, Helbling A, Berchtold E: Immunotherapy with honeybee venom and yellow jacket venom is different regarding efficacy and safety. *J Allergy Clin Immunol* 1992;89:529–535.
- Golden DB, Tracy JM, Freeman TM, Hoffman DR: Negative venom skin test results in patients with histories of systemic reaction to a sting. *J Allergy Clin Immunol* 2003;112:495–498.
- Busse WW, Reed CE, Lichtenstein LM, Reisman RE: Immunotherapy in bee-sting anaphylaxis. Use of honeybee venom. *JAMA* 1975;231:1154–1156.
- Golden DB, Meyers DA, Kagey-Sobotka A, Valentine MD, Lichtenstein LM: Clinical relevance of the venom-specific immunoglobulin G antibody level during immunotherapy. *J Allergy Clin Immunol* 1982;69:489–493.
- Celesnik Smodis N, Silar M, Erzen R, Rijavec M, Kosnik M, Korosec P: Down-regulation of FcεRI-mediated CD63 basophil response during short-term VIT determined venom-nonspecific desensitization. *PLoS One* 2014;9:e94762.
- Uermosi C, Beerli RR, Bauer M, Manolova V, Dietmeier K, Buser RB, Kundig TM, Saudan P, Bachmann MF: Mechanisms of allergen-specific desensitization. *J Allergy Clin Immunol* 2010;126:375–383.

- 32 Schmitz N, Dietmeier K, Bauer M, Maudrich M, Utzinger S, Muntwiler S, Saudan P, Bachmann MF: Displaying Fel d 1 on virus-like particles prevents reactogenicity despite greatly enhanced immunogenicity: a novel therapy for cat allergy. *J Exp Med* 2009;206: 1941–1955.
- 33 Bachmann MF, Kundig TM: Allergen-specific immunotherapy: is it vaccination against toxins after all? *Allergy* 2017;72:13–23.
- 34 Visco V, Dolecek C, Denepoux S, Le Mao J, Guret C, Rousset F, Guinnepain MT, Kraft D, Valenta R, Weyer A, Banchereau J, Labecque S: Human IgG monoclonal antibodies that modulate the binding of specific IgE to birch pollen Bet v 1. *J Immunol* 1996;157:956–962.
- 35 Kemeny DM, MacKenzie-Mills M, Harries MG, Youten LJ, Lessof MH: Antibodies to purified bee venom proteins and peptides. II. A detailed study of changes in IgE and IgG antibodies to individual bee venom antigens. *J Allergy Clin Immunol* 1983;72:376–385.
- 36 Pereira-Santos MC, Baptista AP, Melo A, Alves RR, Soares RS, Pedro E, Pereira-Barbosa M, Victorino RM, Sousa AE: Expansion of circulating Foxp3+ CD25bright CD4+ T cells during specific venom immunotherapy. *Clin Exp Allergy* 2008;38:291–297.
- 37 Blaauw PJ, Smithuis LO: The evaluation of the common diagnostic methods of hypersensitivity for bee and yellow jacket venom by means of an in-hospital insect sting. *J Allergy Clin Immunol* 1985;75:556–562.
- 38 Kampelmacher MJ, van der Zwan JC: Provocation test with a living insect as a diagnostic tool in systemic reactions to bee and wasp venom: a prospective study with emphasis on the clinical aspects. *Clin Allergy* 1987;17:317–327.